

A Guide to Treating Your Condition with

HYFTOR[®] (sirolimus topical gel) 0.2%

Not an actual patient.
Individual results may vary.

Tube not shown
at actual size.

What is HYFTOR[®]?

HYFTOR[®] is a prescription medicine that is used on the skin (topical) to treat adults and children 6 years of age and older with a type of noncancerous tumor called angiofibroma on your face caused by the genetic condition Tuberous Sclerosis Complex.

It is not known if HYFTOR[®] is safe and effective in children under 6 years of age.



Learn more at
HYFTOR.com

IMPORTANT SAFETY INFORMATION

Important: HYFTOR[®] is for use on the skin only (topical use). Do not use HYFTOR[®] in your mouth, eyes, or vagina.

Do not use HYFTOR[®] if you are allergic to sirolimus or any of the other ingredients in HYFTOR[®].

This Guide is intended for US residents only. The information provided is for educational purposes. Please see additional Important Safety Information, including Patient Information and the full Prescribing Information on pages 14 to 23.

Facial Angiofibroma and TSC

Facial angiofibroma is a condition consisting of benign (noncancerous) tumors that appear as pinkish or reddish bumps, usually on the cheeks, nose, and chin. These tumors may be associated with the rare genetic disease Tuberous Sclerosis Complex (TSC). Although these tumors are benign, they may still require treatment.

Facial angiofibroma occurs in approximately 75%-80% of patients with TSC, making it one of the most common skin conditions seen in this population.

Some patients may be successfully treated with invasive procedures (such as surgical removal, laser therapy, or dermabrasion), which may require anesthesia.

About HYFTOR®

HYFTOR® is a clear topical gel that contains sirolimus, an mTOR inhibitor. The mTOR pathway may play a role in the development of TSC. By targeting this pathway, HYFTOR® may help reduce the size and redness of facial angiofibromas.

HYFTOR®
(sirolimus topical gel) 0.2%



Talk to your doctor about your condition and if HYFTOR® is right for you.

Important Safety Information (cont'd)

Before using HYFTOR®, tell your healthcare provider about all of your medical conditions, including if you:

- have a skin infection at the treatment site
- have high cholesterol or high triglycerides (fat or lipids) in your blood are scheduled to receive an immunization (vaccine). You should avoid receiving live vaccines during treatment with HYFTOR®. Vaccines may be less effective during treatment with HYFTOR®.

Please see additional Important Safety Information, including Patient Information and the full Prescribing Information on pages 14 to 23.

Improvement in Facial Angiofibroma Associated with Tuberous Sclerosis Complex at Week 12

HYFTOR® n = 30 patients



Placebo n = 32 patients



- "Improved" or "Markedly Improved"
- "Improved"
- "Markedly Improved"

Before and After Images of Clinical Study Results.



Individual results may vary. JAMA Dermatology. 2018;154(7):781-788. © 2018 American Medical Association. All rights reserved.

- An assessment of "improved" was defined as at least a 50% reduction in size and a 2-level reduction in redness, and an assessment of "markedly improved" was defined as at least a 75% reduction in size and a 3-level reduction in redness
- 62 patients participated in the trial (including 25 patients between 6 and 18 years of age)
- 30 patients applied HYFTOR® and 32 patients applied a placebo (a gel containing no medication) to their faces twice daily for 12 weeks
- Results were reported at the end of 12 weeks. The improvement in size and redness of facial angiofibromas was assessed by the investigator (live assessment) using the subjects' original baseline photographs as reference

Important Safety Information (cont'd)

Before using HYFTOR®, tell your healthcare provider about all of your medical conditions, including if you (cont'd):

- are pregnant or plan to become pregnant. HYFTOR® may harm your unborn baby. You should not become pregnant during treatment with HYFTOR®.
- Females who are able to become pregnant should use effective birth control (contraception) before starting treatment with HYFTOR®, during treatment, and for 12 weeks after your final dose of HYFTOR®. Talk to your healthcare provider about types of birth control that you can use during this time.
- are breastfeeding or plan to breastfeed. It is not known if HYFTOR® passes into your breast milk. You should not breastfeed during treatment with HYFTOR®

The Science Behind HYFTOR[®]

Mechanism of Action

This interview with Dr. Darcy Krueger, Professor of Neurology at Cincinnati Children's hospital, along with an animation, explains how HYFTOR[®] works by regulating cell growth that may cause facial angiofibroma associated with TSC.



WATCH VIDEO



HYFTOR[®]
(sirolimus topical gel) 0.2%

Please see additional Important Safety Information, including Patient Information and the full Prescribing Information on pages 14 to 23.

How to Use HYFTOR®

HYFTOR®
(sirolimus topical gel) 0.2%

Talk to your doctor about your dose and how to apply HYFTOR®. HYFTOR® is only for topical use on the area of the face affected with angiofibroma. The skin being treated with HYFTOR® should not be covered with bandages, dressings, or wraps.



HYFTOR® should be applied to the skin of the face affected with angiofibroma twice daily, in the morning and at bedtime.



Store HYFTOR® refrigerated at 36° to 46°F. Protect from light.



Wash your hands before and after applying HYFTOR®.



Tell your healthcare provider if the treated skin area does not improve within 12 weeks of treatment.




Tube not shown
at actual size.

The maximum recommended daily dosage is:

- Approximately 3/4 inch of gel (600 mg) for pediatric patients 6 to 11 years of age
- Approximately 1 inch of gel (800 mg) for adults and pediatric patients 12 years of age and older

Use the dose depicted below twice daily or as recommended by your doctor.



Sunlight and UV Exposure

Limit your exposure to sunlight and ultraviolet (UV) light during treatment with HYFTOR®. Wear protective clothing and use sunscreen when outdoors.

Important Safety Information (cont'd)

Before using HYFTOR®, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. HYFTOR® may harm your unborn baby. You should not become pregnant during treatment with HYFTOR®.
- Females who are able to become pregnant should use effective birth control (contraception) before starting treatment with HYFTOR®, during treatment, and for 12 weeks after your final dose of HYFTOR®. Talk to your healthcare provider about types of birth control that you can use during this time.

What should I avoid while using HYFTOR®?

Limit your exposure to sunlight and artificial light, such as tanning beds and ultraviolet light therapy, during treatment with HYFTOR®. Wear clothing that covers your skin if you need to go outside. Talk with your healthcare provider about other ways you can protect your skin from the sun.

How to Use

HYFTOR[®]

(sirolimus topical gel) 0.2%



WATCH VIDEO



Not an actual patient. Individual results may vary.

Please see additional Important Safety Information, including Patient Information and the full Prescribing Information on pages 14 to 23.

Side Effects

The most common side effects, occurring in 1% or more of patients treated with HYFTOR®, are:

- Dry skin
- Application site irritation
- Itching
- Acne
- Acne-like rash
- Eye redness
- Skin bleeding
- Skin irritation

These are not all the possible side effect of HYFTOR®. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.



Not an actual patient.
Individual results may vary.

HYFTOR®
(sirolimus topical gel) 0.2%

Please see additional Important Safety Information, including Patient Information and the full Prescribing Information on pages 14 to 23.

Treatment Diary

Check the box after each application.
Write any changes you notice in the Notes section.



Weeks 1-4

Notes

Notes

Notes

Notes

Weeks 5-8

Notes

Notes

Notes

Notes

Weeks 9-12

Notes

Notes

Notes

Notes

Reaching 12 weeks is an important milestone. Please share your progress with your doctor.

Tell your healthcare provider if the treated skin area does not improve within 12 weeks of treatment.

Please see additional Important Safety Information, including Patient Information and the full Prescribing Information on pages 14 to 23.

Discussion Guide to prepare for your next appointment

Preparation Checklist

The questions below can help you think about some of the topics that may come up at your next appointment. Sharing this information may help your healthcare team recommend a treatment plan that is right for you. Bring your answers with you to help guide your conversation with your healthcare provider.



Note to Parents/Caregivers

It can be helpful to discuss these topics with your loved one before their appointment so you can both be prepared.

Question 1 How is your facial angiofibroma? What is your current experience with the condition?

Your answer 1

Question 2 Has anything changed since your last appointment? Has your facial angiofibroma gotten worse or better?

Your answer 2

Question 3 How are you treating your facial angiofibroma? How is it going? Are you still using it?

Your answer 3

Question 4 Are you experiencing any challenges with your current treatment? If so, what are they?

Your answer 4

Question 5 Are you experiencing any side effects?

Your answer 5

Question 6 Is anything else keeping you from continuing your treatment?

Your answer 6

Question 7 Are you having any trouble getting access to your current treatment? Are you having issues with insurance coverage or anything else?

Your answer 7

Question 8 Are you interested in learning about any new treatment options?

Your answer 8

**Contribute to Society by Developing
Critical but Neglected Pharmaceuticals
and Medical Devices**

Nobelpharma

HYFTOR[®]

(sirolimus topical gel) 0.2%

Important Safety Information

What is HYFTOR[®]?

HYFTOR[®] is a prescription medicine that is used on the skin (topical) to treat adults and children 6 years of age and older with a type of noncancerous tumor called angiofibroma on your face caused by the genetic condition tuberous sclerosis.

It is not known if HYFTOR[®] is safe and effective in children under 6 years of age.

Important: HYFTOR[®] is for use on the skin only (topical use). Do not use HYFTOR[®] in your mouth, eyes, or vagina.

Do not use HYFTOR[®] if you are allergic to sirolimus or any of the other ingredients in HYFTOR[®].

Before using HYFTOR[®], tell your healthcare provider about all of your medical conditions, including if you:

- have a skin infection at the treatment site
- have high cholesterol or high triglycerides (fat or lipids) in your blood
- are scheduled to receive an immunization (vaccine). You should avoid receiving live vaccines during treatment with HYFTOR[®]. Vaccines may be less effective during treatment with HYFTOR[®].

- are pregnant or plan to become pregnant. HYFTOR[®] may harm your unborn baby. You should not become pregnant during treatment with HYFTOR[®].
- Females who are able to become pregnant should use effective birth control (contraception) before starting treatment with HYFTOR[®], during treatment, and for 12 weeks after your final dose of HYFTOR[®]. Talk to your healthcare provider about types of birth control that you can use during this time.
- are breastfeeding or plan to breastfeed. It is not known if HYFTOR[®] passes into your breast milk. You should not breastfeed during treatment with HYFTOR[®].

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What should I avoid while using HYFTOR[®]?

Limit your exposure to sunlight and artificial light, such as tanning beds and ultraviolet light therapy, during treatment with HYFTOR[®]. Wear clothing that covers your skin if you need to go outside. Talk with your healthcare provider about other ways you can protect your skin from the sun.

What are possible side effects of HYFTOR[®]? HYFTOR[®] may cause serious side effects, including:

- Allergic reactions. Serious allergic reactions have happened in people who have taken sirolimus by mouth. Stop using HYFTOR® and get medical help right away if you get any of these symptoms: swelling of your face, eyes, or mouth, trouble breathing or wheezing, throat tightness, chest pain or tightness, feeling dizzy or faint, rash or peeling of your skin.
- Serious infections, including infections that can happen when your immune system is weak, have happened in people who have taken sirolimus by mouth. Some people have developed a rare, serious brain infection called progressive multifocal leukoencephalopathy (PML) which can sometimes cause death. Stop using HYFTOR® and call your healthcare provider right away if you get symptoms of an infection including fever or chills.
- Risk of cancer. Lymphoma and other cancers, especially skin cancer, have happened in people who have taken sirolimus by mouth. Talk with your healthcare provider about your risk for cancer if you use HYFTOR®.
- Increased levels of cholesterol and triglycerides (fat or lipids) in the blood have happened in people who have taken sirolimus by mouth. Your healthcare provider will treat you for

high lipid levels, if needed.

- Lung or breathing problems. Lung or breathing problems, including problems that have sometimes caused death, have happened in people who have taken sirolimus by mouth. Stop using HYFTOR® and get medical help right away if you get symptoms such as shortness of breath, new or worsening cough, or chest pain.

The most common side effects of HYFTOR® include dry skin, application site irritation, itching, acne, acne-like rash, eye redness, skin bleeding, and skin irritation.

HYFTOR® may cause fertility problems in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effect of HYFTOR®, Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You are encouraged to report side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088. You may also reach out to Nobelpharma America, LLC at 1-877-375-0825

Please see full Prescribing Information including Patient Information at page 16 to 23

PATIENT INFORMATION HYFTOR® (HYFE TORE) (SIROLIMUS TOPICAL GEL)

Important: HYFTOR is for use on the skin only (topical use). Do not use HYFTOR in your mouth, eyes, or vagina.

What is HYFTOR?

HYFTOR is a prescription medicine that is used on the skin (topical) to treat adults and children 6 years of age and older with a type of noncancerous tumor called angiofibroma on your face caused by the genetic condition tuberous sclerosis. It is not known if HYFTOR is safe and effective in children under 6 years of age.

Do not use HYFTOR if you are allergic to sirolimus or any of the other ingredients in HYFTOR. See the end of this leaflet for a complete list of ingredients in HYFTOR.

Before using HYFTOR, tell your healthcare provider about all of your medical conditions, including if you:

- have a skin infection at the treatment site
- have high cholesterol or high triglycerides (fat or lipids) in your blood
- are scheduled to receive an immunization (vaccine). You should avoid receiving live vaccines during treatment with HYFTOR. Vaccines may be less effective during treatment with HYFTOR.
- are pregnant or plan to become pregnant. HYFTOR may harm your unborn baby. You should not become pregnant during treatment with HYFTOR.
 - » Females who are able to become pregnant should use effective birth control (contraception) before starting treatment with HYFTOR, during treatment, and for 12 weeks after your final dose of HYFTOR. Talk to your healthcare provider about types of birth control that you can use during this time.
- are breastfeeding or plan to breastfeed. It is not known if HYFTOR passes into your breast milk. You should not breastfeed during treatment with HYFTOR.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using HYFTOR with certain medicines may affect each other causing side effects.

How should I use HYFTOR?

Use HYFTOR exactly as your healthcare provider tells you to use it.

- Before you use HYFTOR, your healthcare provider or pharmacist should show you how to correctly measure your dose.
- Wash your hands before and after applying HYFTOR.
- Apply HYFTOR to the skin of the face affected with angiofibroma 2 times a day, in the morning and at bedtime.
- **Do not** cover, wrap, apply dressings, or bandage the skin area treated with HYFTOR.
- Tell your healthcare provider if the treated skin area does not improve within 12 weeks of treatment.

What should I avoid while using HYFTOR?

Limit your exposure to sunlight and ultraviolet light, such as tanning beds and ultraviolet light therapy, during treatment with HYFTOR. Wear clothing that covers your skin if you need to go outside. Talk with your healthcare provider about other ways you can protect your skin from the sun.

What are the possible side effects of HYFTOR?

HYFTOR may cause serious side effects, including:

- **Allergic reactions.** Serious allergic reactions have happened in people who have taken sirolimus by mouth. Stop using HYFTOR and get medical help right away if you get any of these symptoms of a serious allergic reaction:
 - » swelling of your face, eyes, or mouth
 - » chest pain or tightness
 - » trouble breathing or wheezing
 - » feeling dizzy or faint
 - » throat tightness
 - » rash or peeling of your skin
- **Infections.** Serious infections, including infections that can happen when your immune system is weak, have happened in people who have taken sirolimus by mouth. Some people have developed a rare, serious brain infection called progressive multifocal leukoencephalopathy (PML) which can sometimes cause death. Stop using HYFTOR and call your healthcare provider right away if you get symptoms of an infection including fever or chills.
- **Risk of cancer.** Lymphoma and other cancers, especially skin cancer, have happened in

people who have taken sirolimus by mouth. Talk with your healthcare provider about your risk for cancer if you use HYFTOR.

- » **Increased levels of cholesterol and triglycerides (fat or lipids) in the blood** have happened in people who have taken sirolimus by mouth. Your healthcare provider may do blood tests to check you for high lipid levels during treatment with HYFTOR and treat you, if needed.
- **Lung or breathing problems.** Lung or breathing problems, including problems that have sometimes caused death, have happened in people who have taken sirolimus by mouth. Stop using HYFTOR and get medical help right away if you get symptoms such as shortness of breath, new or worsening cough, or chest pain.

The most common side effects of HYFTOR include dry skin, application site irritation, itching, acne, acne-like rash, eye redness, skin bleeding, and skin irritation.

HYFTOR may cause fertility problems in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects of HYFTOR.

Call your doctor for medical advice about side

effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store HYFTOR ?

- Store HYFTOR in the refrigerator between at 36°F to 46°F (2°C to 8°C).
 - Keep HYFTOR out of light.
- Keep HYFTOR and all medicines out of the reach of children.**

General information about the safe and effective use of HYFTOR.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use HYFTOR for a condition for which it was not prescribed. Do not give HYFTOR to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about HYFTOR that is written for health professionals.

What are the ingredients in HYFTOR?

Active ingredient:

Inactive ingredients: alcohol 51%, Carbomer 940, purified water, and trolamine.

Distributed by: Nobelpharma America, LLC
4520 East-West Highway, Suite 400, Bethesda, MD 20814

For more information, go to www.Hyftor.com or call 887-375-0825.

This Patient Information has been approved by the
U.S. Food and Drug Administration.

101-104-02
Issued: 5/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1. INDICATIONS AND USAGE
2. DOSAGE AND ADMINISTRATION
3. DOSAGE FORMS AND STRENGTHS
4. CONTRAINDICATIONS
5. WARNINGS AND PRECAUTIONS
 - 5.1 Hypersensitivity Reactions
 - 5.2 Serious Infection
 - 5.3 Malignancy
 - 5.4 Hyperlipidemia
 - 5.5 Interstitial Lung Disease/Non-Infectious Pneumonitis
 - 5.6 Immunizations
 - 5.7 Embryo-Fetal Toxicity
 - 5.8 Male Infertility
6. ADVERSE REACTIONS
 - 6.1 Clinical Trials Experience
7. DRUG INTERACTIONS
 - 7.1 Effects of Other Drugs on HYFTOR
 - 7.2 Effects of HYFTOR on Other Drugs

8. USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.3 Females and Males of Reproductive Potential
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
11. DESCRIPTION
12. CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.3 Pharmacokinetics
13. NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
14. CLINICAL STUDIES
16. HOW SUPPLIED/STORAGE AND HANDLING
17. PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1. INDICATIONS AND USAGE

HYFTOR is indicated for the treatment of facial angiofibroma associated with tuberous sclerosis in adults and pediatric patients 6 years of age and older.

2. DOSAGE AND ADMINISTRATION

- Complete all age-appropriate vaccinations as recommended by current immunization guidelines prior to HYFTOR initiation [see *Warning and Precautions* (5.6)].
- Apply HYFTOR to the skin of the face affected with angiofibroma twice daily in the morning and at bedtime.
- The maximum recommended daily dosage is:
 - » 600 mg (2 cm) for pediatric patients 6 to 11 years of age
 - » 800 mg (2.5 cm) for adults and pediatric patients 12 years of age and older
- If symptoms do not improve within 12 weeks of treatment, reevaluate the need for continuing HYFTOR.
- Do not use HYFTOR with occlusive dressings.
- For topical use only. Not for oral, ophthalmic, or intravaginal use.

3. DOSAGE FORMS AND STRENGTHS

Topical gel, 0.2%: Each gram contains 2 mg of sirolimus in a colorless and transparent gel in 10-gram tubes.

4. CONTRAINDICATIONS

HYFTOR is contraindicated in patients with a history of hypersensitivity to sirolimus or any other component of HYFTOR. Reactions to sirolimus have included anaphylactic/anaphylactoid reactions, angioedema, exfoliative dermatitis, and hypersensitivity vasculitis [see *Warning and Precautions* (5.1)].

5. WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylactic/anaphylactoid reactions, angioedema, exfoliative dermatitis, and hypersensitivity vasculitis have been associated with the oral administration of sirolimus. The concomitant use of HYFTOR with other drugs known to cause angioedema, such as angiotensin-converting enzyme (ACE) inhibitors, may increase the risk of developing angioedema. Elevated sirolimus levels (with or without concomitant ACE inhibitors) may also potentiate angioedema. Discontinue HYFTOR immediately if symptoms of hypersensitivity occur.

5.2 Serious Infection

Serious infections, including opportunistic

infections, have been reported after oral administration of sirolimus. Cases of progressive multifocal leukoencephalopathy (PML), sometimes fatal have been reported in patients treated with oral sirolimus. Discontinue HYFTOR immediately if symptoms of infection occur.

5.3 Malignancy

Lymphoma and other malignancies, particularly of the skin, have been observed after oral administration of sirolimus. Patients should minimize or avoid exposure to natural or artificial sunlight (tanning beds or UVA/Btreatment) while using HYFTOR. If patients need to be outdoors while using HYFTOR, they should wear protective clothing and discuss other sun protection measures with their physician.

5.4 Hyperlipidemia

Increased serum cholesterol and triglycerides requiring treatment have been observed with oral administration of sirolimus. Monitor for hyperlipidemia during treatment with HYFTOR.

5.5 Interstitial Lung Disease/Non-Infectious Pneumonitis

Cases of interstitial lung disease [ILD] (including pneumonitis, bronchiolitis obliterans organizing pneumonia [BOOP], and pulmonary fibrosis), some fatal, with no identified infectious etiology have occurred in patients receiving oral sirolimus. In some cases, the ILD has resolved upon discontinuation or dosage reduction of oral sirolimus. Discontinue HYFTOR immediately if symptoms of ILD occur.

5.6 Immunizations

During treatment with HYFTOR, vaccinations may be less effective. Complete all age-appropriate vaccinations as recommended by current immunization guidelines prior to initiating treatment with HYFTOR. The use of live vaccines should be avoided during treatment with HYFTOR.

5.7 Embryo-Fetal Toxicity

Based on animal studies and the mechanism of action, oral sirolimus can cause fetal harm when administered to a pregnant woman. In animal studies, oral sirolimus caused embryo-fetal toxicity when administered during the period of organogenesis at maternal exposures that were equal to or less than human exposures at the recommended lowest starting dose. HYFTOR is systemically absorbed after topical administration and may result in fetal exposure. Advise pregnant women of the potential risk to a fetus. Advise female patients of reproductive potential to avoid becoming pregnant. They should use effective contraception prior to, throughout treatment and for 12 weeks after the final dose of HYFTOR

[see Use in Specific Populations (8.1, 8.3), Clinical Pharmacology (12.1, 12.3)].

5.8 Male Infertility

Azoospermia or oligospermia has been observed after oral administration of sirolimus [see Use in Specific Populations (8.3), Nonclinical Toxicology (13.1)]. Sirolimus is an anti-proliferative drug and affects rapidly dividing cells like the germ cells. Advise males that HYFTOR may impair fertility.

6. ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In a randomized, double-blind, vehicle-controlled trial, subjects aged 6 years and older with facial angiofibroma associated with tuberous sclerosis applied HYFTOR twice daily for 12 weeks. A total of 30 subjects were treated with HYFTOR and 32 with the vehicle. The majority of the subjects were female (54.8%). A total of 40.3% were less than 18 years of age.

The most common adverse reactions reported by $\geq 1\%$ of subjects treated with HYFTOR and more frequently than in subjects treated with vehicle are presented in Table 1. Adverse reactions occurred with similar frequency in pediatric subjects 6 years of age and older.

Table 1: Adverse Reactions in $\geq 1\%$ of Subjects Aged 6 Years and Older with Facial Angiofibroma Associated with Tuberous Sclerosis Through Week 12

Preferred Term	HYFTOR N = 30	Vehicle N = 32
Dry skin ^a	12 (40%)	4 (13%)
Application site irritation	11 (37%)	9 (28%)
Pruritus	5 (17%)	4 (13%)
Acne	2 (7%)	0 (0%)
Acneiform dermatitis	1 (3%)	0 (0%)
Ocular hyperemia	1 (3%)	0 (0%)
Skin hemorrhage	1 (3%)	0 (0%)
Skin irritation	1 (3%)	0 (0%)

^a Dry skin includes dry skin and asteatosis

In a 104-week, open-label safety trial, the most common adverse reactions associated with HYFTOR application were application site irritation (31%), dry skin (28%), acne (20%), pruritus (9%), eye irritation (9%), erythema (7%), acneiform dermatitis (6%), contact dermatitis (5%), solar

dermatitis (1%), and photosensitivity reaction (1%). Adverse reactions occurred with similar frequency in adult and pediatric subjects 6 years of age and older.

7. DRUG INTERACTIONS

7.1 Effects of Other Drugs on HYFTOR

Table 2 presents clinically significant drug interactions involving drugs that affect HYFTOR.

Table 2: Effects of CYP3A4 Inhibitors on HYFTOR

Clinical Impact	Concomitant use of HYFTOR with inhibitors of CYP3A4 has the potential to increase the systemic exposure of sirolimus and increase the risk of HYFTOR adverse reactions.
Intervention	Monitor for adverse reactions of HYFTOR.

7.2 Effects of HYFTOR on Other Drugs

Systemic exposure of drugs that are both substrates and inhibitors of CYP3A could be increased with coadministration with HYFTOR. Monitor for adverse reactions of such co-administered drugs

8. USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on animal studies and mechanism of action, oral sirolimus can cause fetal harm when administered to a pregnant woman. HYFTOR is systemically absorbed after topical administration and may result in fetal exposure. The available data from case reports on HYFTOR use in pregnant women are insufficient to evaluate for a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Animal reproduction studies have not been conducted with HYFTOR. In an animal reproduction study, oral administration of 0.5 mg/kg/day sirolimus caused embryo-fetal lethality in pregnant rats when administered during the period of organogenesis (see Data). The available data do not allow the calculation of relevant comparisons between the systemic exposure of sirolimus observed in animal studies to the systemic exposure that would be expected in humans after topical use of HYFTOR.

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

In embryo-fetal development studies in rats, oral administration of sirolimus to pregnant rats during the period of organogenesis (gestation days 6 -15) induced embryo-fetal lethality at 0.5 mg/kg/day, reduced fetal body weight at 1.0 mg/kg/day and caused maternal toxicity at 2.0 mg/kg/day. No treatment related embryo-fetal developmental effects were observed at 0.1 mg/kg/day.

In embryo-fetal development studies in rabbits, oral administration of sirolimus to pregnant rabbits during the period of organogenesis (gestation days 6 -18) induced maternal toxicity (decreased body weight) at 0.05 mg/kg/day, which was associated with embryo-fetal loss or early resorption. No treatment related developmental effects were observed at 0.025 mg/kg/day.

In a pre- and postnatal development toxicity study, oral administration of sirolimus to pregnant rats during gestation and lactation (gestation day 6 through lactation day 20) increased the incidence of dead pups, resulting in reduced live litter size at 0.5 mg/kg/day. No treatment related developmental effects were observed at 0.1 mg/kg/day.

Sirolimus did not cause maternal toxicity or affect developmental parameters in the surviving offspring (morphological development, motor activity, learning, or fertility assessment) at 0.5 mg/kg/day, the highest dose tested.

8.2 Lactation

Risk Summary

There are no available data on the presence of sirolimus in human milk, the effects on the breastfed infant, or the effects on milk production. After oral administration, sirolimus was present in the milk of lactating rats. Because of the potential for serious adverse reactions in the breastfed infant, breastfeeding is not recommended during treatment with HYFTOR.

8.3 Females and Males of Reproductive Potential Contraception

Based on animal studies with oral sirolimus, HYFTOR may cause fetal harm when administered to pregnant women. Females of reproductive potential are recommended to avoid becoming pregnant and to use an effective contraceptive method. Effective contraception should be initiated before HYFTOR therapy and used throughout treatment and for 12 weeks after the final dose of HYFTOR [see *Warnings and Precautions* (5.7), *Use in Specific Populations* (8.1)].

Infertility

Based on clinical findings and findings in animal studies, male and female fertility may be compromised by the treatment with sirolimus [see *Warnings and Precautions* (5.8), *Nonclinical Toxicology* (13.1)]. Ovarian cysts and menstrual disorders (including amenorrhea and menorrhagia) have been reported in females with the use of oral sirolimus. Azoospermia has been reported in males with the use of oral sirolimus and have been reversible upon discontinuation of sirolimus in most cases.

8.4 Pediatric Use

The safety and effectiveness of HYFTOR have been established in pediatric patients aged 6 years and older for the topical treatment of facial angiofibroma associated with tuberous sclerosis. Use of HYFTOR in this age group is supported by data from a randomized, vehicle-controlled, double-blind 12-week trial along with additional efficacy and long-term safety data from a 104-week open label safety trial. A total of 13 pediatric subjects aged 6 years to 17 years received HYFTOR in the Phase 3 clinical trial along with 48 pediatric subjects aged 3 years to 17 years in the 104-week open label safety trial. Adverse reactions occurred with similar frequency in adult and pediatric subjects [see *Adverse Reaction* (6.1), *Clinical Studies* (14)].

The safety and effectiveness of HYFTOR for this indication have not been established in pediatric patients less than 6 years of age.


8.5 Geriatric Use

Clinical studies of HYFTOR did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

11. DESCRIPTION

HYFTOR® (sirolimus topical gel) 0.2% is an mTOR inhibitor immunosuppressant for topical use. Each gram contains 2 mg of sirolimus, which is solubilized in a gel consisting of alcohol 51%, Carbomer 940, purified water, and tromamine.

Chemically, sirolimus is designated as (3*S*,6*R*,7*E*,9*R*,10*R*,12*R*,14*S*,15*E*,17*E*,19*E*,21*S*,23*S*,26*R*,27*R*,34*aS*)-14,21,22,23,24,25,26,27,32,33,34,34a-hexadecahydro-9,27-dihydroxy-3-[(1*R*)-2-[(1*S*,3*R*,4*R*)-4-hydroxy-3-methoxycyclohexyl]-1-methylethyl]-10,21-dimethoxy-6,8,12,14,20,26-hexamethyl-23,27-epoxy-3*H*-pyrido[2,1-*c*][1,4]oxaazacyclohentriacontine-1,5,11,28,29(4*H*,6*H*,31*H*)-pentone. It has the following structural formula:



Sirolimus is a white to off-white powder and is insoluble in water, but freely soluble in chloroform, acetone and acetonitrile. Sirolimus has a molecular formula of $C_{51}H_{79}NO_{13}$ and a molecular weight of 914.19.

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of sirolimus in the treatment of angiofibroma associated with tuberous sclerosis is unknown. Tuberous sclerosis is associated with genetic defects in TSC1 and TSC2 which leads to the constitutive activation of mammalian target of rapamycin (mTOR). Sirolimus inhibits mTOR activation.

12.3 Pharmacokinetics

Absorption

Following 12 weeks of treatment with HYFTOR in adult and pediatric subjects aged 6 years and older, sirolimus blood concentrations ranged from undetectable to 0.50 ng/mL after multiple doses of HYFTOR in the Phase 3 trial. Periodic blood samples were obtained in the 52-week trial and the maximum sirolimus concentration measured at any time in adult subjects was 3.27 ng/mL and the maximum sirolimus concentration measured at any time in pediatric subjects was 1.80 ng/mL.

Distribution

There was no evidence based on blood concentrations that sirolimus accumulates systemically upon topical application in patients with tuberous sclerosis for periods of up to 1 year.

Elimination

Metabolism

Studies evaluating the metabolism of HYFTOR have not been conducted.

Sirolimus is a substrate for both CYP3A4 and P-gp. Sirolimus is extensively metabolized after oral administration in the intestinal wall and liver and undergoes counter-transport from enterocytes of the small intestine into the gut lumen. Inhibitors of CYP3A4 and P-gp increase sirolimus concentrations. Inducers of CYP3A4 and P-gp decrease sirolimus concentrations. Sirolimus is extensively metabolized by O-demethylation and/or hydroxylation. Seven (7) major metabolites, including hydroxy, demethyl, and hydroxydemethyl, are identifiable in whole blood. Some of these metabolites are also detectable in plasma, fecal, and urine samples. Sirolimus is the major component in human whole blood and contributes to more than 90% of the immunosuppressive activity.

Excretion

Studies evaluating the excretion of HYFTOR have not been conducted.

After a single dose of [^{14}C] sirolimus oral solution in healthy volunteers, the majority (91%) of radioactivity was recovered from the feces, and only a minor amount (2.2%) was excreted in urine. The mean $\pm$ SD terminal elimination half-life ($t_{1/2}$) of sirolimus after multiple dosing in stable renal transplant patients was estimated to be about 62 ± 16 hours.

Drug Interaction Studies

Drug interaction studies with HYFTOR have not been conducted. Sirolimus is known to be a substrate for both cytochrome P-450 3A4 (CYP3A4) and p-glycoprotein (P-gp). [see *Drug Interactions* (7.1, 7.2)].

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Oral carcinogenicity studies were conducted in mice and rats. In an oral carcinogenicity study conducted in female mice, sirolimus administered once daily for 86 weeks was associated with a statistically significant increase in malignant lymphoma at all dose levels compared with control animals. In a second oral mouse carcinogenicity study, sirolimus-related hepatocellular adenoma and carcinoma were induced in male mice. In an oral carcinogenicity study conducted in rats with sirolimus there were no significant findings.

Sirolimus was not genotoxic in the in vitro bacterial reverse mutation assay, the Chinese hamster ovary cell chromosomal aberration assay, the mouse lymphoma cell forward mutation assay, or the in vivo mouse micronucleus assay.

Fertility was decreased in both male and female rats following oral administration of sirolimus 2.0 and 0.5 mg/kg, respectively. In male rats, atrophy of testes, epididymides, prostate, seminiferous tubules and/or reduced sperm counts were observed. In female rats, decreased ovarian and uterine weights and decreased implantation were observed. Testicular tubular degeneration was also seen in a 4-week intravenous study of sirolimus in monkeys at 0.1 mg/kg.

14. CLINICAL STUDIES

A single, randomized, double-blind, vehicle-controlled, multi-center, Phase 3 trial was conducted in Japan to evaluate HYFTOR for the treatment of adults and pediatric patients 6 years of age and older with facial angiofibroma associated with tuberous sclerosis (NCT02635789). A total of 62 Japanese subjects with 3 or more angiofibromas (≥ 2 mm in diameter with redness in each) on the face were enrolled in this trial. Overall, 28 subjects (45%) were male and 34 (55%) were female. A total of 25 subjects (40%) were between 6 and <18 years of age. In this trial, subjects applied either HYFTOR or vehicle twice daily to the skin of their face affected with angiofibroma for 12 weeks.

The efficacy was assessed by the investigator (live assessment) based on the composite improvement from baseline in size and redness of facial angiofibroma, using subjects' baseline photographs as reference. The proportion of subjects assessed as 'Improved' or 'Markedly Improved' at Week 12 is presented in Table 3. An assessment of 'Improved' was defined as at least a 50% reduction in the size and a 2-level reduction in redness and an assessment of 'Markedly Improved' was defined as at least a 75% reduction in the size and a 3-level reduction in redness

Table 3: Improvement in Facial Angiofibroma Associated with Tuberous Sclerosis in Patients Aged 6 Years and Older at Week 12

Proportion of Subjects Assessed by the Investigator as:	HYFTOR N=30	Vehicle N=32
'Improved' or 'Markedly Improved'	23%	6%
'Improved'	13%	3%
'Markedly Improved'	10%	3%

16. HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

HYFTOR® (sirolimus topical gel) 0.2% is a colorless and transparent gel supplied as 10 g in aluminum tubes (NDC 73683-101-10). Each gram contains 2 mg of sirolimus.

Storage and Handling

Store refrigerated at 2° to 8°C (36° to 46°F). Protect from light.

17. PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Hypersensitivity

Inform patients that oral sirolimus has been associated with hypersensitivity reactions, including anaphylactic/anaphylactoid reactions, angioedema, exfoliative dermatitis, and hypersensitivity vasculitis. Advise patients to discontinue HYFTOR immediately and seek medical attention if symptoms occur [see *Warnings and Precautions* (5.1)].

Serious Infections

Inform patients that oral sirolimus has been associated with increased susceptibility to infections, including opportunistic infections and latent viral infections, such as progressive multifocal leukoencephalopathy (PML). Advise patients to discontinue HYFTOR immediately and seek medical attention if symptoms occur [see *Warnings and Precautions* (5.2)].

Malignancy

Inform patients that oral sirolimus has been associated with malignancy, including lymphoma and skin cancer. Advise patients to minimize or avoid exposure to natural or artificial sunlight (tanning beds or UVA/B treatment), and to use protective measures if exposure cannot be avoided, while using HYFTOR [see *Warnings and Precautions* (5.3)].

Hyperlipidemia

Inform patients that oral sirolimus has been associated with increased serum cholesterol and triglycerides requiring treatment and that periodic laboratory monitoring may be needed [see *Warnings and Precautions* (5.4)].

Interstitial Lung Disease

Inform patients that oral sirolimus has been associated with interstitial lung disease [ILD] sometimes fatal, with no identified infectious etiology. Advise patients to discontinue HYFTOR immediately and seek medical attention if symptoms (e.g. shortness of breath) occur [see *Warnings and Precautions* (5.5)].

Immunization

Inform patients that during treatment with HYFTOR, vaccinations may be less effective. Instruct patients that vaccination with live vaccines should be avoided during treatment with HYFTOR and to inform the healthcare practitioner that they are

using HYFTOR prior to a potential vaccination [see *Warnings and Precautions* (5.6)].

Pregnancy

HYFTOR may cause fetal harm if used during pregnancy. Advise female patients of reproductive potential to avoid becoming pregnant and to use effective contraception prior to, throughout treatment, and for 12 weeks after the final dose of HYFTOR. Advise pregnant women of the potential risk to a fetus [see *Warnings and Precautions* (5.7), *Use in Specific Populations* (8.1, 8.3)].

Lactation

Advise lactating women that breastfeeding is not

recommended during treatment with HYFTOR [see *Use in Specific Populations* (8.2)].

Infertility

Inform male and female patients that HYFTOR may impair fertility [see *Warnings and Precautions* (5.8), *Use in Specific Populations* (8.3), *Nonclinical Toxicology* (13.1)].

Administration Information

Inform patients that the skin being treated with HYFTOR should not be covered with bandages, dressings or wraps [see *Dosage and Administration* (2)]

Distributed by:



Nobelpharma America, LLC
4520 East-West Highway, Suite 400
Bethesda, MD 20814

Nobelpharma America Co-Pay Program

Pay as Little as \$0*



The Co-Pay Card may help eligible, commercially-insured US patients pay for certain out-of-pocket, drug-related expenses.

The Co-Pay card is not insurance, but it may compliment your insurance. Your specialty pharmacy will call you to coordinate payment and delivery of your prescription. Please inquire about your eligibility for the Co-Pay program. If you have any questions, please don't hesitate to contact Nobelpharma directly at **1-877-649-3867** or visit nobelpharmaconnect.com.

Other Nobelpharma Connect Programs

- **Patient Assistance Program**
For eligible un-insured patients
- **Bridge Program**
For facing a brief insurance lapse
- **Quick Start Program**
For eligible new patients facing insurance delays

*Available to patients with commercial insurance coverage for Nobelpharma America pharmaceutical product who meet eligibility criteria. This Co-Pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. Please call us at 1-877-649-3867 for additional information.



Please visit NobelpharmaConnect.com for more information

Nobelpharma